

# Buckeye® Lamb Milk Replacer

# 110925

TYPE OF FEED Complete

FORM OF FEED: Powder

PACKAGING: 25 lb paper bags

SPECIES: Lambs (may also be fed to goat kids and deer fawn)

AGES: newborn



---

## FEATURES:

- Provides the entire nutritional requirements of the baby lamb from its second day of life through weaning
- May be fed on a warm restricted basis or fed cold ad lib in a cafeteria type feeder
- Lactose level is reduced as lambs cannot digest the milk sugar at significant levels
- Contains no medication
- The protein in this product is derived solely from milk
- Acidified to help inhibit development of enteric pathogens in the stomach which will increase lamb survival and improve growth rates pre-weaning

## GUARANTEED ANALYSIS

Crude Protein, min.	23.00%
Crude Fat, min.	30.00%
Crude Fiber, max.	0.20%
Calcium, min.	0.45%
Calcium, max.	0.95%
Phosphorus, min	0.60%
Selenium, min	0.30 ppm
Vitamin A, min.	20,000 IU/lb
Vitamin D, min.	5,000 IU/lb
Vitamin E, min.	50 IU/lb

INGREDIENTS: Dried Whey Protein Concentrate, Dried Whey, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Whey Product, Lecithin, DL Methionine, L-Lysine, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Vitamin B-12 Supplement, Folic Acid, Choline Chloride, Riboflavin Supplement, Niacin Supplement, Calcium Pantothenate, Thiamine Monoitrate, [ Sodium Propionate, Citric Acid, Potassium Sorbate, Calcium Propionate(Preservatives)], Ferrous Sulfate, Cobalt Sulfate, Zinc Sulfate, Manganese Sulfate, Magnesium Oxide, Ethylenediamine Dihydriodide, Sodium Silico Aluminate, Sodium Selenite, Artificial Flavor.

FEEDING DIRECTIONS: See Second Page

# Buckeye® Lamb Milk Replacer

## FEEDING DIRECTIONS:

**FEEDING DIRECTIONS (Lambs):** Mix at the rate of 2 lb. per gallon of water or 0.25 lb.\* of Buckeye® Lamb Milk Replacer with 16 ounces (1 pint) of water.

1. Ensure that the baby lamb receives colostrum during the first 12 hours of life.
2. For hand feeding with a nipple bottle or restricted feeding, offer warm at 100 degrees Fahrenheit. It is recommended to offer 3 to 6 feedings per day of warm milk during the first two weeks of life and thereafter 2 feedings per day. Never offer more than ½ pint per feeding.
3. For best results, mix the lamb milk replacer with cold water and offer ad libitum to lambs through a suitable nipple feeder. Milk should be available at all times. Reconstituted Lamb Milk Replacer should be changed and equipment cleaned every 48 hours or less.
4. Provide Buckeye® 18% Lamb Starter in a lamb creep at 10 to 14 days of age and restrict hay intake to the amount the lambs will consume in one hour.
5. Wean lambs at a minimum of 30 days of age, weighing at least 25 pounds.

**(Goat Kids)** Mix at the rate of 1.5 lb. Per gallon of water (.2 lb. of powder per pint of water).

1. Goat kid can be fed on a restricted basis or offered ad-libitum through a suitable feeder. In all cases, ensure kids receive colostrum during the first 24 hours after partition. For the first few days, feed a limited quantity of a maximum of 1 pint per feeding up to 3 to 4 times a day. As the kids get older, reduce the number of feedings per day.
2. Start goat kids on Buckeye® 16% Goat Feed at 5 to 8 days of age. At 4 weeks of age limited amounts of excellent quality hay should be introduced to the kids. If coccidiosis is a problem with the young kids, they may be fed Buckeye® 18% Calf Starter medicated with Deccox.  
\*One level plastic measuring cup enclosed in the bag equals .25 lb. of dry Buckeye® Lamb Milk Replacer.