

Advance Kid Milk Replacer

110432

TYPE OF FEED Complete

FORM OF FEED: Powder

PACKAGING: 25 lb paper bags

SPECIES: Lambs (may also be fed to goat kids and deer fawn)

AGES: newborn



GUARANTEED ANALYSIS

Crude Protein, min.	23.00%
Crude Fat, min.	30.00%
Crude Fiber, max.	0.20%
Calcium, min.	0.45%
Calcium, max.	0.95%
Phosphorus, min	0.60%
Selenium, min	0.30 ppm
Vitamin A, min.	20,000 IU/lb
Vitamin D, min.	5,000 IU/lb
Vitamin E, min.	50 IU/lb

INGREDIENTS: Dried Whey Protein Concentrate, Dried Whey, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Whey Product, Lecithin, DL Methionine, L-Lysine, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Vitamin B-12 Supplement, Folic Acid, Choline Chloride, Riboflavin Supplement, Niacin Supplement, Calcium Pantothenate, Thiamine Mononitrate, [Sodium Propionate, Citric Acid, Potassium Sorbate, Calcium Propionate(Preservatives)], Ferrous Sulfate, Cobalt Sulfate, Zinc Sulfate, Manganese Sulfate, Magnesium Oxide, Ethylenediamine Dihydriodide, Sodium Silico Aluminate, Sodium Selenite, Artificial Flavor.

FEEDING DIRECTIONS: See Second Page

Buckeye® Lamb Milk Replacer

FEEDING DIRECTIONS:

FEEDING DIRECTIONS (Lambs): Mix at the rate of 2 lb. per gallon of water or 0.25 lb.* of Buckeye® Lamb Milk Replacer with 16 ounces (1 pint) of water.

1. Ensure that the baby lamb receives colostrum during the first 12 hours of life.

2. For hand feeding with a nipple bottle or restricted feeding, offer warm at 100 degrees Fahrenheit. It is recommended to offer 3 to 6 feedings per day of warm milk during the first two weeks of life and thereafter 2 feedings per day, Never offer more than ½ pint per feeding.
3. For best results, mix the lamb milk replacer with cold water and offer ad libitum to lambs through a suitable nipple feeder. Milk should be available at all times. Reconstituted Lamb Milk Replacer should be changed and equipment cleaned every 48 hours or less.
4. Provide Buckeye® 18% Lamb Starter in a lamb creep at 10 to 14 days of age and restrict hay intake to the amount the lambs will consume in one hour.
5. Wean lambs at a minimum of 30 days of age, weighing at least 25 pounds.

(Goat Kids) Mix at the rate of 1.5 lb. Per gallon of water (.2 lb. of powder per pint of water).

1. Goat kid can be fed on a restricted basis or offered ad-libitum through a suitable feeder. In all cases, ensure kids receive colostrum during the first 24 hours after partition. For the first few days, feed a limited quantity of a maximum of 1 pint per feeding up to 3 to 4 times a day. As the kids get older, reduce the number of feedings per day.

2. Start goat kids on Buckeye® 16% Goat Feed at 5 to 8 days of age. At 4 weeks of age limited amounts of excellent quality hay should be introduced to the kids. If coccidiosis is a problem with the young kids, they may be fed Buckeye® 18% Calf Starter medicated with Deccox.

*One level plastic measuring cup enclosed in the bag equals .25 lb. of dry Buckeye® Lamb Milk Replacer.